

On the homotopy type of a cofibred category

MATIAS L. DEL HOYO

Departamento de Matematica
FCEyN, Universidad de Buenos Aires
Buenos Aires, Argentina.

Abstract

In this paper we describe two ways on which cofibred categories give rise to bisimplicial sets. The *fibred nerve* is a natural extension of Segal's classical nerve of a category, and it constitutes an alternative simplicial description of the homotopy type of the total category. If the fibration is splitting, then one can construct the *cleaved nerve*, a smaller variant which emerges from a distinguished closed cleavage. We interpret some classical theorems by Thomason and Quillen in terms of our constructions, and use the fibred and cleaved nerve to establish new results on homotopy and homology of small categories.

Introduction

The classifying space functor associates to every small category C a topological space BC , namely the geometric realization of its nerve [Se]. The classical homotopy theory of categories is lifted from spaces by using this functor. For instance, a *weak equivalence* between small categories is a map $f : C \rightarrow C'$ such that Bf is a homotopy equivalence.

A fundamental fact concerning this construction is that for every space X there is a small category C such that X and BC have the same weak homotopy type (cf. [Il, VI,3.3.1], see also [dH]). This way small categories constitute models for homotopy types, and one seeks to characterize the discrete invariants of X in terms of its underlying category C .

It is natural to expect that a small category C endowed with extra structure would give rise to a space BC equipped with some additional data. That is our motivation for introducing the *fibred nerve* and the *cleaved nerve*. These are bisimplicial sets with the homotopy type of the total category of a Grothendieck fibration, and constitute combinatorial description that preserve in some sense the fibred structure.

By a Grothendieck fibration, or just a *fibration*, we mean what is usually called a *cofibred category*. We adopt this notation for simplicity, and to emphasize the analogy with the topological case. Other notions of fibrations between small categories have been studied, for instance, in [Mi, Ev].

Grothendieck fibrations have played an important role in homotopy theory. Among others, they were used by Thomason to describe homotopy colimits of small categories [Th], and Quillen's Theorems A and B – that lead to long exact sequences of higher K-theory groups – may be stated in terms of Grothendieck fibrations [Qu]. We believe that the nerve constructions studied here will help in further applications, such as explicit constructions of $K(G, n)$ categories and Postnikov towers in Cat .

Organization

Section 1 deals with preliminaries. We fix some notations and recall some results about the classifying space functor and a key proposition on simplicial sets (1.2.1). The reader is referred to [Qu] for an introduction to homotopy of small categories, and to [GJ] for a comprehensive treatment of bisimplicial objects.

2000 *Mathematics Subject Classification*: 18D30; 18G30; 55U35.

Key words: Cofibred category; Classifying space; Nerve.

The principal reference on Grothendieck fibrations is [SGA1, VI]. A more recent one is [Bo]. In section 2 we set the definitions, recall some facts about fibrations and develop some others which will be needed later, such as the correspondence 2.2.3.

In section 3 we introduce both the fiber and the cleaved nerve, in the same fashion as the classical nerve is defined. We establish some fundamental facts (cf. 3.1.2, 3.2.2) and prove that for a splitting fibration the two constructions yield the same homotopy type (cf. 3.2.3).

We prove that the fibred nerve is homotopy equivalent to the classic nerve in section 4 (cf. 4.1.3). From these we derive the original and the relative versions of Quillen's theorem A. In addition, we show how to recover the classic nerve of a splitting fibration using the codiagonal construction over a bisimplicial set (cf. 4.3.3).

Last section summarizes applications and relations between the fibred nerve and some other constructions.

- The cleaved nerve and Bousfield-Kan construction for homotopy colimits are related in 5.1.1. We derive Thomason's theorem on homotopy colimits of small categories as a consequence.
- We develop a Leray-Serre style spectral sequence (cf. 5.2.1) relating the homology groups of the base, the several fibers and the total category. We deduce as a corollary a homology version of Quillen's Theorem A (cf. 5.2.3).
- We introduce Quillen fibrations, which are families of categories with the same homotopy type, and show that Quillen's Theorem B might be interpreted as the following conceptual fact: the fibred classifying space functor maps Quillen fibrations into quasifibrations (5.3.1).
- Finally, we associate to a category endowed with a group action a splitting fibration, and prove that its cleaved nerve is a twisted cartesian product as defined in [Ma].

Acknowledgements

I would like to thank Gabriel Minian, my advisor. His several suggestions and remarks were essential in the development and revision of this work. I also thank to Fernando Cukierman and Eduardo Dubuc for many stimulating talks. Lastly, I thank to CONICET for the financial support.

1 Preliminaries

We denote by $\mathbf{Cat}$, $\mathbf{SSet}$ and $\mathbf{Top}$ the categories of small categories, simplicial sets and topological spaces, respectively. If C is a small category, then we denote by $\mathbf{ob}(C)$ its set of objects and by $\mathbf{fl}(C)$ its set of arrows. As usual, with Δ we denote the category of finite ordinals, and with $[n]$ we mean the ordinal of $n + 1$ elements. We write I for the simplicial set represented by $[1]$. Sometimes $[n]$ will be regarded as a category in the usual sense.

1.1 About homotopy of small categories

Given C a small category, its *nerve* NC is the simplicial set whose n -simplices are the chains

$$\underline{c} = (c_0 \rightarrow c_1 \rightarrow \dots \rightarrow c_n)$$

of n composable arrows in C , and its *classifying space* BC is the geometric realization of its nerve, namely $BC = |NC|$. It is a CW-complex with one n -cell for each chain of n composable arrows in C which does not involve an identity [Se].

A functor $f : C \rightarrow C'$ in $\mathbf{Cat}$ is a *weak equivalence* if Bf is a homotopy equivalence in $\mathbf{Top}$, and a small category C is *contractible* if BC is so. From the homeomorphism $B(C \times I) \cong BC \times BI$ it follows

that a functor $C \times I \rightarrow C'$ induces a continuous map $BC \times [0, 1] \rightarrow BC'$ and therefore a natural transformation $h : f \Rightarrow g : C \rightarrow C'$ yields a homotopy $Bh : Bf \Rightarrow Bg : BC \rightarrow BC'$. This leads to the following results [Qu].

1.1.1 Lemma. If a functor admits an adjoint, then it is a weak equivalence.

1.1.2 Lemma. A category having an initial or final object is contractible.

It is well known that given a commutative triangle in Top , if two of the three involved arrows are weak homotopy equivalences, then so does the third. It follows immediately that the same statement holds for weak equivalences in SSet and weak equivalences in Cat . We will refer to this fact as the *3for2-property*.

1.2 About bisimplicial sets

The nerve of a category is a simplicial set. We shall extend this concept by constructing the fibred nerve of a fibration, which is a bisimplicial set. A *bisimplicial set* is a functor $K : \Delta^\circ \times \Delta^\circ \rightarrow \text{Set}$, where Δ° denotes the opposite category of Δ . A bisimplicial set K can be regarded as a family of sets $\{K_{m,n}\}_{m,n \geq 0}$ equipped with horizontal and vertical faces and degeneracies operators satisfying some simplicial identities [GJ]. We denote by bSSet the category of bisimplicial sets and morphisms between them.

A bisimplicial set is the same as a simplicial object in SSet . Given K a bisimplicial set, let $\{m \mapsto K_{m,n}\}$ be the n -th *vertical* simplicial set, which is obtained from K by setting the second coordinate equal to n . The m -th *horizontal* simplicial set $\{n \mapsto K_{m,n}\}$ is defined analogously. We denote by $d(K)$ the *diagonal* of K , namely the simplicial set which is the composition of K with the diagonal functor $\Delta^\circ \rightarrow \Delta^\circ \times \Delta^\circ$.

We define the *geometric realization* of K as the space $|d(K)|$, which is naturally homeomorphic to the spaces obtained by first realizing on one direction and then on the other [Qu, p.10].

$$|n \mapsto |m \mapsto K_{m,n}|| \cong |d(K)| \cong |m \mapsto |n \mapsto K_{m,n}||$$

If $f : K \rightarrow L$ is a map of bisimplicial sets, then we say that it is a *weak equivalence* if its geometric realization $f_* : |d(K)| \rightarrow |d(L)|$ is a homotopy equivalence. The following is a very useful criterion to establish when a map is a weak equivalence (see e.g. [BK, XII,2.3] or [GJ, IV,1.9]).

1.2.1 Proposition. Let $f : X \rightarrow Y$ be a map in bSSet such that for all n the induced map $f_* : \{m \mapsto X_{m,n}\} \rightarrow \{m \mapsto Y_{m,n}\}$ is a weak equivalence in SSet . Then f is a weak equivalence.

2 Fibrations

If $u : A \rightarrow B$ is a map between small categories, we say that $f \in \text{fl}(A)$ is *over* $\phi \in \text{fl}(B)$ if $u(f) = \phi$, and we say that $f \in \text{fl}(A)$ is *over* $b \in \text{ob}(B)$ if $u(f) = \text{id}_b$. Given $b \in \text{ob}(B)$, the *fiber* u_b is the subcategory of A of arrows over b , and the *homotopy fiber* u/b is the category whose objects are pairs (a, ϕ) , $a \in \text{ob}(A)$ and $\phi : u(a) \rightarrow b \in \text{fl}(B)$, and whose arrows $f : (a, \phi) \rightarrow (a', \phi')$ are maps $f : a \rightarrow a'$ in A such that $\phi' u(f) = \phi$. By an abuse of notation we shall write A_b and A/b instead of u_b and u/b . Note that there is a canonical fully faithful inclusion $A_b \rightarrow A/b$, defined by $a \mapsto (a, \text{id}_{u(a)})$.

2.1 Basic definitions and examples

Let $p : E \rightarrow B$ a map between small categories. An arrow $f : e \rightarrow e'$ in E is said to be *cartesian* if it satisfies the following universal property: for all $g : e \rightarrow e''$ over $p(f)$ there is a unique $h : e' \rightarrow e''$

over $p(e')$ such that $hf = g$.

$$\begin{array}{ccc} e & \xrightarrow{f} & e' \\ & \searrow \forall g & \downarrow \exists! h \\ & & e'' \end{array}$$

$$p(e) \xrightarrow{p(f)} p(e')$$

A map $p : E \rightarrow B$ is a *prefibration* if for all e object of E and all $\phi : p(e) \rightarrow b$ arrow of B there is a cartesian arrow $f : e \rightarrow e'$ over ϕ . It is not hard to see that $p : E \rightarrow B$ is a prefibration if and only if the inclusion $E_b \rightarrow E/b$ of the actual fiber into the homotopy fiber admits a left adjoint for all object b in B . Therefore, if $p : E \rightarrow B$ is a prefibration then the inclusion $E_b \rightarrow E/b$ is a weak equivalence for all b (cf. 1.1.1).

A prefibration $p : E \rightarrow B$ is called a *fibration* if cartesian arrows are closed under composition. We say that B is the *base category* and that E is the *total category* of the fibration.

Examples.

- The projection $\pi : F \times B \rightarrow B$ is a fibration, since the arrows $(id, \phi) \in \text{fl}(F \times B)$ are cartesian.
- Given B a small category and $F : B \rightarrow \text{Cat}$ a functor, the projection $F \rtimes B \rightarrow B$ is a fibration, whose fibers are the values of F . Here $F \rtimes B$ denotes the *Grothendieck construction* over F (see e.g. section 5.1).
- We denote by B^I the category of functors $I \rightarrow B$. Its objects are the arrows of B and its maps $(u, v) : f \rightarrow g$ are the commutative squares $vf = gu$ in B . The functor $\text{cod} : B^I \rightarrow B$ which assigns to each arrow its codomain is a fibration. The fibers of cod are the *slice categories* B/b .
- If $A \subset B$ is a *coideal* (cf. [He]), then the inclusion $A \rightarrow B$ is a fibration, whose fibers are either $\emptyset$ or pt , the final object of Cat .
- Given $p : E \rightarrow B$, an isomorphism $f \in \text{fl}(E)$ is always cartesian. Thus, a functor between groupoids that is onto on arrows is a fibration.

The cartesian arrows in a fibration satisfy the following stronger universal property (cf. [Bo]).

2.1.1 Lemma. Let $p : E \rightarrow B$ be a fibration and $f : e \rightarrow e'$ a cartesian arrow in E . Given $g : e \rightarrow e''$ such that $p(g) = \phi p(f)$ for some $\phi : p(e') \rightarrow p(e'')$, there exists a unique arrow $h : e' \rightarrow e''$ over ϕ satisfying $hf = g$.

$$\begin{array}{ccc} e & \xrightarrow{f} & e' \\ & \searrow \forall g & \downarrow \exists! h \\ & & e'' \end{array}$$

$$p(e) \xrightarrow{p(f)} p(e') \xrightarrow{\phi} p(e'')$$

Moreover, h is cartesian if and only if g is so.

Given a prefibration $p : E \rightarrow B$, a *cleavage* Σ is a choice of cartesian arrows. More precisely, a cleavage is a subset $\Sigma \subset \text{fl}(E)$ whose elements are cartesian arrows and such that for all $e \in \text{ob}(E)$ and $\phi : p(e) \rightarrow b \in \text{fl}(B)$ there exists a unique arrow $\Sigma_{e,\phi} : e \rightarrow e'$ in Σ over ϕ .

The cleavage Σ is said to be *normal* if it contains the identities, and is said to be *closed* if it is closed under composition. Every prefibration admits a normal cleavage, but not every prefibration admits a closed one. A fibration which admits a closed cleavage is called a *splitting fibration*.

Example. Let E, B be groups, regarded as categories with a single object, and let $p : E \rightarrow B$ be a map between them. Then every map of E is cartesian as it is an isomorphism. It follows that p is a fibration if and only if p is an epimorphism of groups. A cleavage Σ for p is a set-theoretic section for p . The cleavage is normal if Σ preserves the neutral element, and the cleavage is closed if it is a morphism of groups. This example shows in particular that “only a few” fibrations are splitting.

From here on we will assume that all the cleavages are normal. The following lemma, whose proof is straight-forward, gives an alternative description of closed cleavages.

2.1.2 Lemma. A cleavage Σ is closed if and only if $f \in \Sigma$ and $f'f \in \Sigma$ imply that $f' \in \Sigma$ for all pair f, f' of composable arrows of E .

Next we discuss two notions of morphism between fibrations, and describe the corresponding categories.

Given $\xi = (p : E \rightarrow B)$ and $\xi' = (p' : E' \rightarrow B')$ fibrations, we define a *fibred map* $\xi \rightarrow \xi'$ as a pair $f : E \rightarrow E', g : B \rightarrow B'$ of maps in Cat such that f preserves cartesian arrows and $p'f = gp$.

$$\begin{array}{ccc} E & \xrightarrow{f} & E' \\ \downarrow p & & \downarrow p' \\ B & \xrightarrow{g} & B' \end{array}$$

We denote by $\text{hom}_{\text{fib}}(\xi, \xi')$ the set of fibred maps $\xi \rightarrow \xi'$, and by Fib the category of fibrations and fibred maps between them.

Now suppose that cleavages Σ and Σ' of ξ and ξ' are given. A *cleaved map* $(\xi, \Sigma) \rightarrow (\xi', \Sigma')$ is a fibred map $(f, g) : \xi \rightarrow \xi'$ such that $f(\Sigma) \subset \Sigma'$. By $\text{hom}_{\text{cliv}}((\xi, \Sigma), (\xi', \Sigma'))$ we mean the set of cleaved maps $(\xi, \Sigma) \rightarrow (\xi', \Sigma')$, and by Cliv the category of pairs (ξ, Σ) and cleaved maps.

Finally, we denote by Esc the full subcategory of Cliv whose objects are the pairs (ξ, Σ) with Σ a closed cleavage of ξ .

We have the following diagram, where the first is a full inclusion and the arrow $\text{Cliv} \rightarrow \text{Fib}$ is the forgetful functor $(\xi, \Sigma) \mapsto \xi$.

$$\text{Esc} \subset \text{Cliv} \rightarrow \text{Fib} \subset \text{Cat}^I$$

2.2 Fibration associated to a map

Given $u : A \rightarrow B$ a map between small categories, we define the *mapping category* E^u as the fiber product $A \times_B B^I$ over u and $\text{dom} : B^I \rightarrow B$ in Cat . The objects of E^u are pairs $(a, u(a) \rightarrow b)$, with a an object of A and $u(a) \rightarrow b$ an arrow of B , and the arrows are pairs (f, g) which induce a commutative square in B .

The functor u factors through E^u as πi , where i is the inclusion $a \mapsto (a, \text{id}_{u(a)})$, and π is the projection $(a, u(a) \rightarrow b) \mapsto b$.

$$\begin{array}{ccccc} A & \xrightarrow{i} & E^u & \xrightarrow{\pi} & B \\ & \searrow u & & & \end{array}$$

The functor i is fully faithful and admits a right adjoint, the retraction $r : E^u \rightarrow A$, which maps $(a, u(a) \rightarrow b)$ into a . This implies the following (cf. 1.1.1).

2.2.1 Lemma. The map $i : A \rightarrow E^u$ is a weak equivalence.

The functor π is a fibration. The set $\Sigma^u \subset \text{fl}(E^u)$ of arrows whose first coordinate is an identity

$$\Sigma^u = \{(\text{id}_a, \phi) : (a, u(a) \rightarrow b) \rightarrow (a, u(a) \rightarrow b')\},$$

is a closed cleavage for π , so it is a splitting fibration. We say that $\pi : E^u \rightarrow B$ is the *fibration associated to u* , and we endow it with the cleavage Σ^u . Note that if b is an object of B , then the fiber E_b^u of π is isomorphic to the homotopy fiber A/b of u .

Except in very special situations, the retraction $r : E^u \rightarrow A$ does not commute with the projections, namely (r, id_B) is not a map in Cat^I . We shall describe how to replace r by others well-behaved retractions when the map u is already a fibration.

Let $p : E \rightarrow B$ be a fibration, and let $\pi : E^p \rightarrow B$ be its associated fibration. We say that a map $s : E^p \rightarrow E$ is *good* if $si = id_E$, $ps = \pi$ and s preserves cartesian arrows.

$$\begin{array}{ccc} E & \xrightleftharpoons[i]{s} & E^p \\ p \searrow & & \swarrow \pi \\ & B & \end{array}$$

If s is good, then $(s, id_B) : E^p \rightarrow E$ is a fibred map.

2.2.2 Lemma. A good map s is a weak equivalence, and it induces a weak equivalence $s : E_b^p \rightarrow E_b$ for all object b in B .

Proof. The first statement holds by 2.2.1 since s is left inverse to i . About the second, note that under the isomorphism $E_b^p \cong E/b$ the induced map $E_b^p \rightarrow E_b$ identifies with a left inverse to the inclusion $E_b \rightarrow E/b$, which is a weak equivalence indeed. $\square$

2.2.3 Proposition. Given $p : E \rightarrow B$ a fibration, there is a 1-1 correspondence between normal cleavages of E and good maps $s : E^p \rightarrow E$.

Proof. Let $s : E^p \rightarrow E$ be a good map. For each $e \in \text{ob}(E)$ and $\phi : p(e) \rightarrow b \in \text{fl}(B)$ the arrow $(id_e, \phi) : (e, id : p(e) \rightarrow p(e)) \rightarrow (e, \phi : p(e) \rightarrow b)$ is cartesian in E^p . Therefore, $s(id_e, \phi)$ is a cartesian arrow of E over ϕ with domain $s(i(e)) = e$. It follows that the family $\Sigma = \{s(id_e, \phi)\}_{e, \phi}$ is a cleavage of E , and it is normal because $s(id_e, id_{p(e)}) = s(i(id_e)) = id_e$.

Conversely, if Σ is a normal cleavage of E , then we shall construct a good map $s = s(\Sigma) : E^p \rightarrow E$ as follows. An object $(e, \phi : p(e) \rightarrow b)$ in E^p is mapped by s into the codomain of $\Sigma_{e, \phi} \in \text{ob}(E)$. An arrow $(\alpha, \beta) : (e, \phi : p(e) \rightarrow b) \rightarrow (e', \phi' : p(e') \rightarrow b')$ of E^p is mapped by s into the unique arrow over β which makes the following diagram commutative.

$$\begin{array}{ccccc} e & \xrightarrow{\alpha} & e' & \xrightarrow{\Sigma_{e', \phi'}} & \\ \searrow \Sigma_{e, \phi} & & & & \\ & s(e) & \xrightarrow{s(\alpha, \beta)} & s(e') & \\ & & & & \\ p(e) & \xrightarrow{p(\alpha)} & p(e') & \xrightarrow{\phi'} & \\ \searrow \phi & & & & \\ & b & \xrightarrow{\beta} & b' & \end{array}$$

The uniqueness of $s(\alpha, \beta)$ follows from 2.1.1. It also follows from 2.1.1 that s preserves cartesian arrows. As it respects identities and compositions, s is indeed a functor, and $ps = \pi$ by construction. The map s defined this way is a retraction for $i : E \rightarrow E^p$ because Σ is normal.

It is straightforward to check that these procedures are mutually inverse. $\square$

If E is endowed with a cleavage Σ and $s : E^p \rightarrow E$ is a good map such that $s(\Sigma^u) \subset \Sigma$, then we say that s is *very good*. If s is very good, then $(s, id_B) : (E^p, \Sigma^p) \rightarrow (E, \Sigma)$ is a cleaved map.

2.2.4 Corollary. If s and Σ are related as in 2.2.3, then Σ is closed if and only if the map s is very good.

Proof. Let Σ be a closed cleavage and s its induced good map. If (id_e, β) is an arrow in Σ^u , then the diagram of above gives $s(id_e, \beta)\Sigma_{e, \phi} = \Sigma_{e, \beta\phi}$. It follows from 2.1.2 that $s(id_e, \beta) \in \Sigma$ and hence the map s is very good.

On the other hand, given Σ a cleavage which is not closed, by 2.1.2 one can find f and f' cartesian arrows of E such that $f' = gf$ with $g \notin \Sigma$. Since $g = s(id, p(g))$ it follows that s is not very good. $\square$

3 Bisimplicial sets from fibrations

3.1 Fibred nerve

For $m, n \geq 0$ let $\square_{m,n}$ denotes the fibration $pr_2 : [m] \times [n] \rightarrow [n]$. These are the fibrations which play the role of simplices in Fib . They define a covariant functor $\square : \Delta \times \Delta \rightarrow \text{Fib}$.

Given $\xi = (p : E \rightarrow B)$ a fibration, we define the *fibred nerve* of ξ as the bisimplicial set $N_f \xi$ whose m, n -simplices are given by

$$N_f \xi_{m,n} = \text{hom}_{\text{fib}}(\square_{m,n}, \xi).$$

We define the *fibred classifying space* $B_f \xi$ as the geometric realization $|d(N_f \xi)|$ of the fibred nerve. These constructions are functorial. For short, we shall write $N_f E$ and $B_f E$ instead of $N_f \xi$ and $B_f \xi$.

The fibred nerve extends the classical nerve in the sense that there exists a natural isomorphism

$$d(N_f(id_B)) = d(N_f B) \cong NB.$$

A m, n -simplex of $N_f E$ consists of a pair $s = (s_0, s_1)$, where $s_0 : [m] \times [n] \rightarrow E$ and $s_1 : [n] \rightarrow B$ are such that the induced square commutes. We say that $s_1 \in NB_n$ is the *base* of the simplex s , and that $s_0|_{pr_2^{-1}(0)} \in (NE_{b_0})_m$ is the *mast* of s . Of course, s_0 completely determines s .

We visualize s as an array of arrows of E going down and right. The horizontal arrows are cartesian and the vertical arrows are over identities.

$$\begin{array}{ccccccc} e_{00} & \longrightarrow & e_{01} & \longrightarrow & \dots & \longrightarrow & e_{0n} \\ & & \downarrow & & & & \downarrow \\ e_{10} & \longrightarrow & e_{11} & \longrightarrow & \dots & \longrightarrow & e_{1n} \\ & & \downarrow & & & & \downarrow \\ \vdots & & \downarrow & & \dots & & \downarrow \\ & & \downarrow & & & & \downarrow \\ e_{m0} & \longrightarrow & e_{m1} & \longrightarrow & \dots & \longrightarrow & e_{mn} \\ & & & & & & \\ b_0 & \longrightarrow & b_1 & \longrightarrow & \dots & \longrightarrow & b_n \end{array}$$

Next technical result plays a key role hereafter. Fix $\underline{b} \in NB_n$, and let $N_f E_{\underline{b}}$ be the simplicial set whose simplices are those of $N_f E$ with base $\underline{b}$, with faces and degeneracies in the vertical direction.

3.1.1 Lemma. The map $\mu : N_f E_{\underline{b}} \rightarrow NE_{b_0}$ which gives to each simplex s its mast is a weak equivalence of simplicial sets.

Proof. We choose a cleavage Σ and construct a homotopy inverse $\nu : NE_{b_0} \rightarrow N_f E_{\underline{b}}$ for μ as follows. The map ν associates to a simplex $\underline{a}$ the unique simplex $s = \nu(\underline{a})$ with mast $\underline{a}$ and base $\underline{b}$ and such that $s_0(i \rightarrow i, j \rightarrow j+1) \in \Sigma$ for all i, j . It is clear that $\mu\nu = id$. We shall describe a simplicial homotopy

$$h : N_f E_{\underline{b}} \times I \rightarrow N_f E_{\underline{b}}$$

between $\nu\mu$ and id , which induces a continuous homotopy $|h|$ and completes the proof.

We have that $(N_f E_{\underline{b}} \times I)_m = (N_f E_{\underline{b}})_m \times I_m$, and that $I_m = \{t : [m] \rightarrow [1]\}$. Given $(s, t) \in (N_f E_{\underline{b}} \times I)_m$ we define $h(s, t)$ as the unique m -simplex of $N_f E_{\underline{b}}$ with the same mast as s and such that

$$h(s, t)_0(i \rightarrow i, j \rightarrow j + 1) = \begin{cases} s_0(i \rightarrow i, j \rightarrow j + 1) & \text{if } t(i) = 0 \\ v\mu(s)_0(i \rightarrow i, j \rightarrow j + 1) & \text{if } t(i) = 1 \end{cases}$$

It is easy to see that h defined as above is a simplicial map, that $h(s, 0) = s$ and that $h(s, 1) = v\mu(s)$. $\square$

The main feature of the fibred nerve is that it satisfies the following homotopy preserving property.

3.1.2 Proposition. Let $\xi = (p : E \rightarrow B)$ and $\xi' = (p' : E' \rightarrow B)$ be fibrations, and let $f : E \rightarrow E'$ be a fibred map over B (i.e. $(f, id_B) \in \text{hom}_{fib}(\xi, \xi')$). If $f : E_b \rightarrow E'_b$ is a weak equivalence for all object b of B , then $f_* : N_f E \rightarrow N_f E'$ is a weak equivalence.

Proof. By proposition 1.2.1 it suffices to prove that the map $f_* : \{m \mapsto N_f E_{m,n}\} \rightarrow \{m \mapsto N_f E'_{m,n}\}$ is a weak equivalence for each n . Faces and degeneracies in direction m preserve the base of a simplex, thus we have decompositions

$$\{m \mapsto N_f E_{m,n}\} = \coprod_{\underline{b}=(b_0 \rightarrow \dots \rightarrow b_n)} N_f E_{\underline{b}} \quad \text{and} \quad \{m \mapsto N_f E'_{m,n}\} = \coprod_{\underline{b}=(b_0 \rightarrow \dots \rightarrow b_n)} N_f E'_{\underline{b}}.$$

Moreover, f_* also preserves the base of a simplex, and therefore it can be written as the coproduct of the maps $f_* : N_f E_{\underline{b}} \rightarrow N_f E'_{\underline{b}}$. Now consider the following commutative square.

$$\begin{array}{ccc} N_f E_{\underline{b}} & \xrightarrow{f_*} & N_f E'_{\underline{b}} \\ \downarrow \mu & & \downarrow \mu \\ N E_{b_0} & \xrightarrow{f_*} & N E'_{b_0} \end{array}$$

The vertical maps are weak equivalences by 3.1.1, and the bottom one is so by hypothesis. It follows from the 3for2 property that the upper one is also a weak equivalence and thus the proposition. $\square$

3.2 Cleaved nerve

The fibration $\square_{m,n}$ is splitting, since its unique cleavage $\Sigma = \{(id, \alpha)\}$ is closed. We consider $\square_{m,n}$ as equipped with this cleavage, and we obtain a covariant functor $\square : \Delta \times \Delta \rightarrow \text{Esc} \subset \text{Cliv}$.

Given $\xi = (p : E \rightarrow B)$ a fibration endowed with a cleavage Σ , we define the *cleaved nerve of (ξ, Σ)* as the bisimplicial set $N_c(\xi, \Sigma)$ whose m, n -simplices are given by

$$N_c(\xi, \Sigma)_{m,n} = \text{hom}_{cliv}(\square_{m,n}, (\xi, \Sigma))$$

We define the *cleaved classifying space* of $B_c(\xi, \Sigma)$ as the geometric realization $|d(N_c(\xi, \Sigma))|$ of the cleaved nerve. These constructions are functorial. As before, we shall write $N_c E$ and $B_c E$ instead of $N_c(\xi, \Sigma)$ and $B_c(\xi, \Sigma)$ when there is no place to confusion.

The cleaved nerve extends the classical nerve in the sense that there is a natural isomorphism

$$d(N_c(id_B)) = d(N_c B) \cong NB,$$

where $id : B \rightarrow B$ is equipped with the cleavage $\Sigma = \text{fl}(B)$.

Note that, if we forget the cleavage Σ , then we can form the fibred nerve $N_f E$ and there is a natural inclusion in bSSet

$$i : N_c E \rightarrow N_f E.$$

3.2.1 Lemma. Let $\xi = (p : E \rightarrow B)$ be a fibration with cleavage Σ . If s and s' are simplices in $N_c E$ with the same base and the same mast, then $s = s'$. If Σ is closed, then for all $\underline{b} \in NB_n$ and $\underline{a} \in (NE_{b_0})_m$ there exists a unique m, n -simplex $s \in N_c E$ with base $\underline{b}$ and mast $\underline{a}$.

Proof. Note that $s_0(i \rightarrow i, 0 \rightarrow 1) = s'_0(i \rightarrow i, 0 \rightarrow 1)$ since they are arrows in Σ over $b_0 \rightarrow b_1$ with the same domain. We see that $s_0(i \rightarrow i, j \rightarrow j+1) = s'_0(i \rightarrow i, j \rightarrow j+1)$ by iterating this argument. Finally, $s_0(i \rightarrow i+1, j \rightarrow j) = s'_0(i \rightarrow i+1, j \rightarrow j)$ by the universal property of cartesian arrows. This proves the first assertion.

It is not hard to see that there exists a unique simplex $s \in N_f E$ with base $\underline{b}$, mast $\underline{a}$, and such that $s_0(i \rightarrow i, j \rightarrow j+1) \in \Sigma$ for all i, j . If Σ is closed then $s_0(i \rightarrow i, j \rightarrow k) \in \Sigma$ for all i, j, k and thus s is in $N_c E$ and the second statement holds. $\square$

If the fibration is splitting, then $N_c E$ satisfies a homotopy preserving property analogous to 3.1.2.

3.2.2 Proposition. Let $\xi = (p : E \rightarrow B)$ and $\xi' = (p' : E' \rightarrow B)$ be splitting fibrations with closed cleavages Σ and Σ' , and let $f : E \rightarrow E'$ be a cleaved map over B (i.e. $(f, id_B) \in \text{hom}_{cliv}((\xi, \Sigma), (\xi', \Sigma'))$). If $f : E_b \rightarrow E'_b$ is a weak equivalence for all object b of B then $f_* : N_c E \rightarrow N_c E'$ is a weak equivalence.

Proof. It is analogous to that of 3.1.2, using the restriction $\mu : N_c E_{\underline{b}} \rightarrow NE_{b_0}$, which is also a weak equivalence by 3.2.1 – actually, it is an isomorphism. $\square$

The following result asserts that the cleaved nerve suffices to describe the homotopy type of the fibred nerve when the cleavage is closed.

3.2.3 Theorem. If $\xi = (p : E \rightarrow B)$ is a splitting fibration with closed cleavage Σ , then the inclusion $i : N_c E \rightarrow N_f E$ is a weak equivalence.

Proof. Again by proposition 1.2.1, we only must show that for each n the inclusion induces a weak equivalence $i_* : \{m \mapsto N_c E_{m,n}\} \rightarrow \{m \mapsto N_f E_{m,n}\}$. Fixed n , the map i_* can be written as the coproduct of


$$i_* : N_c E_{\underline{b}} \rightarrow N_f E_{\underline{b}}$$

where $\underline{b}$ runs over all n -simplices of NB . The composition $\mu i_* : N_c E_{\underline{b}} \rightarrow NE_{b_0}$ is an isomorphism by 3.2.1. It follows by 3.1.1 and the 3for2-property that i_* is a weak equivalence and thus the proposition. $\square$

If the cleavage Σ is not closed, then $N_c E$ and $N_f E$ do not necessarily have the same homotopy type. Let us illustrate this with an example.

Example. Let E be the category obtained from the ordinal $[3]$ by formally inverting the arrow $2 \rightarrow 3$. Note that E has an initial element and hence BE is contractible (cf. 1.1.2). We shall see E as the total category of a fibration endowed with a cleavage Σ in such a way that $N_c E$ is not contractible. Since $d(N_f E)$ and NE have the same homotopy type (see 4.1.3), we conclude that in this example the inclusion $i : N_c E \rightarrow N_f E$ is not a weak equivalence.

Let $B = [2]$ and let $p : E \rightarrow B$ be the surjection which takes twice the value 2. Clearly it is a fibration. Let Σ be the normal cleavage which contains the arrow $0 \rightarrow 3$.



By simple inspection one deduces that if a simplex $s \in N_c E$ is not contained in the fiber E_2 , then its mast must be trivial. Since a simplex in $N_c E$ is determined by its mast and its base (cf. 3.2.1), it follows that $0 \in N_c E_{00}$ is face only of $0 \rightarrow 1, 0 \rightarrow 3 \in N_c E_{01}$ and some degenerate simplices. Thus, the loop $0 \rightarrow 1 \rightarrow 2 \rightarrow 3 \leftarrow 0$ gives a non-trivial element of $\pi_1(B_c E, 0)$ and therefore $B_c E$ is not contractible.

4 Relation with the classic nerve

4.1 The main result

Let $\xi = (p : E \rightarrow B)$ be a fibration, and let $s = (s_0, s_1)$ be an element of $N_f E_{n,n}$. The composition $s_0 \circ \text{diag} : [n] \rightarrow E, i \mapsto s_0(i, i)$ gives a n -simplex of NE , which we denote by $k(s)$. This way we get a natural map of simplicial sets

$$k : d(N_f E) \rightarrow NE$$

and its geometric realization $k_* : B_f E \rightarrow BE$. We shall see that it is a weak equivalence, so the fibred nerve becomes an alternative model for the homotopy type of E .

We prove that k is a weak equivalence first for splitting fibrations and then for any fibration.

4.1.1 Proposition. Let $\xi = (p : E \rightarrow B)$ be a splitting fibration, with closed cleavage Σ . Then the map $k|_{d(N_c E)} = ki : d(N_c E) \rightarrow NE$ is a weak equivalence.

Proof. (Compare with [Th, 1.2]) Since 2.2.4 we know that the cleavage Σ induces a very good map $s : E^p \rightarrow E$ and hence a commutative square by the naturality of k .

$$\begin{array}{ccc} d(N_c(E^p)) & \xrightarrow{ki} & N(E^p) \\ \downarrow s_* & & \downarrow s_* \\ d(N_c E) & \xrightarrow{ki} & NE \end{array}$$

In this square the vertical arrows are weak equivalences (cf. 2.2.2, 3.2.2), so in order to prove that the bottom arrow is a weak equivalence, by the 3for2-property it only remains to show that the upper arrow is so. To do that, we define a map $l : d(N_c E^p) \rightarrow NE^p$, prove that there is a simplicial homotopy $ki \cong l$, and prove that l is a weak equivalence.

A simplex $s = (s_0, s_1)$ of $N_c E_{n,n}^p$ is uniquely determined by its mast and its base (cf. 3.2.1), so it consists of the following data

$$s = (e_0 \rightarrow e_1 \rightarrow \dots \rightarrow e_n, p(e_n) \rightarrow b_0 \rightarrow b_1 \rightarrow \dots \rightarrow b_n)$$

Note that $s_0(i, j) = (e_i, p(e_i) \rightarrow b_j)$, where $p(e_i) \rightarrow b_j$ is the arrow induced by s . With these notations,

$$ki(s) = (e_0, p(e_0) \rightarrow b_0) \rightarrow (e_1, p(e_1) \rightarrow b_1) \rightarrow \dots \rightarrow (e_n, p(e_n) \rightarrow b_n).$$

We define $l : d(N_c E^p) \rightarrow NE^p$ by

$$l(s) = (e_0, p(e_0) \rightarrow p(e_n)) \rightarrow (e_1, p(e_1) \rightarrow p(e_n)) \rightarrow \dots \rightarrow (e_n, p(e_n) \rightarrow p(e_n)).$$

The homotopy $h : ki \cong l$ is the map $d(N_c E^p) \times I \rightarrow NE^p$ which in a n -simplex (s, t) is given by

$$h(s, t) = (e_0, p(e_0) \rightarrow p(e_n)) \rightarrow \dots \rightarrow (e_{i-1}, p(e_{i-1}) \rightarrow p(e_n)) \rightarrow (e_i, p(e_i) \rightarrow b_i) \rightarrow \dots \rightarrow (e_n, p(e_n) \rightarrow b_n)$$

where $i = \min\{i : t(i) = 1\}$ if $t \neq 0$ and $i = n + 1$ if $t = 0$. One verifies that h is a map, that $h(s, 0) = l(s)$ and that $h(s, 1) = k(s) = ki(s)$.

Finally, let us prove that l is a weak equivalence. We regard NE^p as a bisimplicial set constant in direction n , so $NE_{m,n}^p = NE_m^p$. The map l , defined above, is the diagonalization of a bisimplicial map $L : N_c E^p \rightarrow NE^p$. The m -th component $L_{m,-}$ of L decomposes as the coproduct

$$\coprod_{e_0 \rightarrow \dots \rightarrow e_m} N(p(e_m)/B) \rightarrow \coprod_{e_0 \rightarrow \dots \rightarrow e_m} \text{pt}$$

which is a weak equivalence because $p(e_m)/B$ has an initial element and therefore is contractible (1.1.2). The map L is a weak equivalence by 1.2.1 and thus the result. $\square$

4.1.2 Corollary. If $\xi = (p : E \rightarrow B)$ is a splitting fibration, then $k : d(N_f E) \rightarrow NE$ is a weak equivalence.

Proof. Fix a closed cleavage Σ and then use 3.2.3 and 4.1.1. □

Now we extend 4.1.2 to a non-necessarily splitting fibration $\xi = (p : E \rightarrow B)$.


4.1.3 Theorem. If $\xi = (p : E \rightarrow B)$ is a fibration, then the map $k : d(N_f E) \rightarrow NE$ is a weak equivalence.

Proof. Let Σ be a cleavage of ξ . The good map $s : E^p \rightarrow E$ induced by Σ (cf. 2.2.3) gives a commutative square

$$\begin{array}{ccc} d(N_f E^p) & \xrightarrow{k} & NE^p \\ s \downarrow & & \downarrow s \\ d(N_f E) & \xrightarrow{k} & NE \end{array}$$

Since the fibration $E^p \rightarrow B$ is always splitting, it follows from 4.1.2 that the upper arrow is a weak equivalence. The vertical arrows are also weak equivalences (cf. 2.2.2, 3.1.2) and then the result follows from the 3for2-property. □

Example. The surjection $s : [2] \rightarrow [1]$ which takes twice the value 1 is a fibration. Down below we show the spaces $B_f E$ and BE . The map k is in this case the obvious inclusion.



Even when this example is quite simple, it is useful to understand some of the differences between the two constructions. Many of the diagonal arrows in the total category do not provide relevant homotopy information, and the fibred nerve omits them.

The cleaved nerve is smaller than the fibred nerve, and therefore a more effective codification of the homotopy type of the total category. On the other hand, it only works when the fibration is splitting, while the fibred nerve is useful for any fibration.

4.2 Quillen's Theorem A and its relative version

Quillen's Theorem A states sufficient conditions for a functor to be a weak equivalence. It was proved to be very useful not only in the work of Quillen but also in many other situations. We derive it here from our framework.

The well-behaviour of fibred nerve with respect to homotopy (cf. 3.1.2) together with theorem 4.1.3 gives the following result.

4.2.1 Proposition. If $f : E \rightarrow E'$ is a fibred map over B such that $f : E_b \rightarrow E'_b$ is a weak equivalence for all object b of B , then f is a weak equivalence.

We deduce both Theorem A and its relative version from last proposition.

4.2.2 Corollary (Relative Quillen's Theorem A). Let $u : A \rightarrow B$ and $u' : A' \rightarrow B$ be small categories over B . If $f : A \rightarrow A'$ is a map over B such that the induced map $A/b \rightarrow A'/b$ is a weak equivalence for all $b \in \text{ob}(B)$, then f is a weak equivalence.

Proof. Consider the following commutative square of categories over B , where E^u and $E^{u'}$ are the associated fibrations to u and u' , and the bottom arrow is induced by f in a natural way.

$$\begin{array}{ccc} A & \xrightarrow{f} & A' \\ i \downarrow & & \downarrow i \\ E^u & \xrightarrow{f_*} & E^{u'} \end{array}$$

Since the actual fiber E_b^u identifies with the homotopy fiber A/b , last proposition asserts that the bottom arrow is a weak equivalence, and the vertical ones are also weak equivalences by 2.2.1. The result follows from this and the 3for2-property. $\square$

4.2.3 Corollary (Quillen's Theorem A). A map $u : A \rightarrow B$ between small categories whose homotopy fibers A/b are contractible is a weak equivalence.

Proof. Take $u' = id_B$ in the relative version. $\square$

4.3 Fibred nerve, cleaved nerve and the codiagonal construction

Given K a bisimplicial set, its *codiagonal* (or *bar construction*) $\nabla(K)$ is the simplicial set whose n -simplices are

$$\nabla(K)_n = \{(x_0, x_1, \dots, x_n) : x_i \in K_{i, n-i}, d_0^h x_i = d_{i+1}^v x_{i+1} \text{ for } 0 \leq i < n\}$$

and whose faces and degeneracies are $d_i(x_0, \dots, x_n) = (d_i^h x_0, d_{i-1}^h x_1, \dots, d_1^h x_{i-1}, d_i^v x_{i+1}, \dots, d_i^v x_n)$ and $s_j(x_0, \dots, x_n) = (s_j^h x_0, s_{j-1}^h x_1, \dots, s_0^h x_j, s_j^v x_{j+1}, \dots, s_j^v x_n)$. There is a natural weak equivalence $\theta : d(K) \rightarrow \nabla(K)$ defined as follows, where x is a n -simplex of $d(K)$ (cf. [CR])

$$\theta(x) = ((d_1^v)^n x, (d_2^v)^{n-1} d_0^h x, \dots, (d_{i+1}^v)^{n-i} (d_0^h)^i x, \dots, (d_0^h)^n x).$$

In the case of the fibred nerve, both the codiagonal $\nabla N_f E$ and the map θ can be described in terms of *singular functors* of fibrations. We shall give these description, and prove that for a splitting fibration there is an isomorphism between the codiagonal of the cleaved nerve and the classic nerve of the total category.

Let T_n be the full subcategory of $[n] \times [n]$ whose objects are the pairs (i, j) satisfying $i \leq j$. The restriction $pr_2|_{T_n} : T_n \rightarrow [n]$ is a fibration as one can easily check. This way we get a covariant functor $T : \Delta \rightarrow \text{Fib}$, $[n] \mapsto T_n$, as the restriction of $\square \circ \text{diag}$.

4.3.1 Proposition. Let $\xi = (p : E \rightarrow B)$ be a fibration. Then there is a canonical isomorphism of simplicial sets

$$\text{hom}_{\text{fib}}(T_n, \xi) \cong (\nabla N_f E)_n$$

where the left member is the "singular functor" induced by $T : \Delta \rightarrow \text{Fib}$. Under this isomorphisms, the map θ identifies with the restriction $s \mapsto s|_{T_n}$.

Proof. For $k = 0, \dots, n$ let $\alpha^k = (\alpha_0^k, \alpha_1^k) : \square_{k, n-k} \rightarrow T_n$ be the fibred map satisfying $\alpha_0^k(i, j) = (i, j+k)$ for all (i, j) . Given $x \in \text{hom}_{\text{fib}}(T_n, \xi)$ note that the composition $x\alpha^k : \square_{k, n-k} \rightarrow \xi$ determines an element in $N_f E_{k, n-k}$. The conditions $d_0^h(x\alpha^k) = d_{k+1}^v(x\alpha^{k+1})$ hold and thus we can define a simplex $\lambda(x) \in (\nabla N_f E)_n$ by

$$\lambda(x) = (x\alpha^0, x\alpha^1, \dots, x\alpha^n)$$

This λ is clearly injective because one can recover x from the compositions $x\alpha^k$. On the other hand, the compatibility conditions among the coordinates of a simplex $y \in \nabla N_f E$ insures that it comes from a map $x : T_n \rightarrow \xi$, which proves that λ is surjective and thus the result. $\square$

4.3.2 Proposition. Let $\xi = (p : E \rightarrow B)$ be a fibration with cleavage Σ . Then there is a canonical isomorphism of simplicial sets

$$\mathrm{hom}_{cliv}(T_n, (\xi, \Sigma)) \cong (\nabla N_c E)_n$$

Given $\xi = (p : E \rightarrow B)$ a fibration endowed with a cleavage Σ , we have the following diagram of simplicial sets, where $\bar{k}$ is defined below.

$$\begin{array}{ccc} dN_c E & \xrightarrow{i} & dN_f E \\ \downarrow \theta & & \downarrow \theta \\ \nabla N_c E & \xrightarrow{i} & \nabla N_f E \end{array} \quad \begin{array}{c} \nearrow k \\ \nearrow \bar{k} \end{array} \quad NE$$

The following shows how to recover the classic nerve of the total category of a splitting fibration from the cleaved nerve.

Proof. The proof is similar to that of lemma 3.2.1. To see that $\bar{k}i$ is injective, consider a simplex $x \in NE_n$, view $x : [n] \rightarrow E$ as defined over the diagonal of $\square_{n,n}$ and note that an extension $s : T_n \rightarrow E$ of x is necessarily unique: The horizontal arrows must belong to the cleavage, and the vertical ones are uniquely determined by the universal property of cartesian arrows.

If Σ is closed, then the unique functor $s : T_n \rightarrow E$ such that $s \circ \text{diag} = x$ and $s((i, j) \rightarrow (i, j + 1)) \in \Sigma$ determines a cleaved map $T_n \rightarrow E$ and hence a simplex $s \in (\nabla_{\mathcal{C}} E)_n$ satisfying $\bar{k}_i(s) = x$. Thus the surjectivity. $\square$

5 Other examples and applications

5.1 Homotopy colimits

Bousfield and Kan established the following explicit construction for homotopy colimits in the category of simplicial sets. Given $Z : I \rightarrow \mathbf{SSet}$, let $hc(Z)$ be the bisimplicial set whose m, n -simplices are

$$hc(Z)_{m,n} = \coprod_{i_0 \rightarrow \dots \rightarrow i_n} Z(i_0)_m$$

where the coproduct runs over all simplices of dimension n of NI , and faces and degeneracies are defined in the obvious way. It turns out that the diagonal complex $d(hc(Z))$ is isomorphic to $hocolim(Z)$ [BK, XII.5.2], or more precisely they are homotopic if we assume that $hocolim(Z)$ is defined “up to homotopy”.

Using Bousfield-Kan description, Thomason construct homotopy colimits of small categories in terms of the Grothendieck construction [Th]. We recall Grothendieck construction over a functor $F : B \rightarrow \mathbf{Cat}$, compare the Bousfield-Kan construction with the cleaved nerve and derive Thomason's theorem from this.

Given $F : B \rightarrow \text{Cat}$ a diagram of small categories, its *Grothendieck construction* is a splitting fibration $F \rtimes B \rightarrow B$ whose fibers are the values of F . The objects of the total category $F \rtimes B$ are pairs (x, b) with b an object of B and x an object of $F(b)$. An arrow $(f, \phi) : (x, b) \rightarrow (x', b')$ in $F \rtimes B$ is a pair $\phi : b \rightarrow b'$, $f : F(\phi)(x) \rightarrow x'$. Composition is given by $(\psi, g) \circ (\phi, f) = (\psi\phi, gF(\psi)(f))$. The map $F \rtimes B \rightarrow B$ is the projection, and the arrows (id, ϕ) form a distinguished closed cleavage.

5.1.1 Theorem. Given $F : B \rightarrow \text{Cat}$, there is an isomorphism $N_c(F \rtimes B) \xrightarrow{\sim} hc(NF)$ between the cleaved nerve of the Grothendieck construction of F and the Bousfield-Kan construction for homotopy colimits.

Proof. The isomorphism $N_c(F \rtimes B) \rightarrow hc(NF)$ maps a m, n -simplex s of $N_c(F \rtimes B)$ to the element $\underline{a}$ in the summand indexed by $\underline{b}$, where $\underline{a}$ is the mast of s and $\underline{b}$ is the base of s . This is indeed a morphism of bisimplicial sets, and it is invertible because of 3.2.1. $\square$

If E is any fibration, then one can define a function $N_f E \rightarrow hc(NF)$ in a similar fashion. However, this function is not a morphism in general, as it does not respect the 0-th face operator.

5.1.2 Corollary (Thomason's theorem). The homotopy colimit of a functor $F : B \rightarrow \text{Cat}$ has the homotopy type of the Grothendieck construction $F \rtimes B$.

Proof. It is a consequence of 4.1.1 and 5.1.1. $\square$

5.2 Spectral sequence of a fibration

A bisimplicial set gives rise to a bisimplicial abelian group and hence to a bicomplex. In this section we study the spectral sequence associated to the bicomplex induced by the fibred nerve. We recall some definitions on homology of categories from [Qu]. Then we describe how a fibration gives rise to a pseudofunctor, and define the modules $H_m(F)$. Finally we state and prove theorem 5.2.1 and derive a homology version of Quillen's Theorem A as a corollary.

Given a small category C , a *module over C* is a functor $A : C \rightarrow \text{Ab}$, where Ab denotes the category of abelian groups. The m -th *homology group of C with coefficients in a module A* is defined as the m -th left derived functor of $\text{colim} : \text{Ab}^C \rightarrow \text{Ab}$.

$$H_m(C, A) = \text{colim}_C^m A$$

The groups $H_m(C, A)$ can be computed as the homology of the following simplicial abelian group

$$C_m(C, A) = \bigoplus_{c_0 \rightarrow \dots \rightarrow c_m} A(c_0)$$

and, in the case that A is morphism inverting, they agree with the homology of the classifying space BC with local coefficients induced by A . We write $H_m(C)$ instead of $H_m(C, A)$ when A is the constant functor $\mathbb{Z}$. It follows that

$$H_m(C) = H_m(BC)$$

where the right side denotes the singular homology of the space BC .

Let $\xi = (p : E \rightarrow B)$ be a fibration, and let Σ be a cleavage of ξ . For each arrow $\phi : b \rightarrow b'$ in B a *base-change functor* $\phi_* : E_b \rightarrow E_{b'}$ is defined as follows: If e is an object of E_b , then $\phi_*(e)$ is the codomain of $\Sigma_{e, \phi}$, and if $f : e \rightarrow e'$ is an arrow of E_b , then $\phi_*(f)$ is the unique arrow in $E_{b'}$ such that $\phi_*(f) \circ \Sigma_{e, \phi} = \Sigma_{e', \phi} \circ f$.

Of course, ϕ_* depends on the cleavage, but different cleavages give rise to naturally isomorphic base-change functors, as it follows from the universal property of cartesian arrows. In the same

fashion, given ϕ, ψ composable arrows of B , there is a natural isomorphism $\psi_*\phi_* \Rightarrow (\psi\phi)_*$. The set of data

$$b \mapsto E_b \quad \phi \mapsto \phi_* \quad \psi_*\phi_* \Rightarrow (\psi\phi)_*$$

defines a *pseudofunctor* $B \dashrightarrow \text{Cat}$ (cf. [SGA1]). Note that if Σ is closed then the isomorphisms $\psi_*\phi_* \Rightarrow (\psi\phi)_*$ are identities and one has a true functor $B \rightarrow \text{Cat}$.

Given $m \geq 0$, let $H_m(F) : B \rightarrow \text{Ab}$ be the functor which assigns to each $b \in B$ the group $H_m(E_b)$, and to each arrow $\phi : b \rightarrow b'$ the map induced by ϕ_* . Since isomorphic functors yields homotopic maps between the classifying spaces, the module $H_m(F)$ is well defined (i.e. is a functor) and does not depend on the cleavage Σ .

5.2.1 Theorem. There is a spectral sequence $\{X_{m,n}^r\}$ which converges to the homology of the total category E and whose second sheet consists of the homology of the base with coefficients in the homology of the fibers.

$$X_{m,n}^2 = H_n(B, H_m(F)) \Rightarrow H_{m+n}(E)$$

Proof. From the bisimplicial set $N_f E$ we construct the free bisimplicial abelian group $\mathbb{Z}N_f E$, and the bicomplex $C_f E$, whose m, n -th group equals that of $\mathbb{Z}N_f E$ and whose horizontal and vertical differential maps are the alternate sum of the horizontal and vertical faces, respectively. We have that

$$C_f E_{m,n} = \bigoplus_{\underline{b}=(b_0 \rightarrow \dots \rightarrow b_n)} \mathbb{Z}[(N_f E_{\underline{b}})_m]$$

where $(N_f E_{\underline{b}})_m$ is the set of m, n -simplices of $N_f E$ with base $\underline{b}$. Filtering the bicomplex $C_f E$ in the horizontal direction gives a spectral sequence

$$H_n(H_m(C_f E)) \Rightarrow H_{m+n}(\text{Tot}(C_f E)).$$

The first sheet of this spectral sequence is obtained by computing the vertical homology (m -direction) of $C_f E$. In degree m, n this is equal to

$$H_m(C_f E)_{m,n} = \bigoplus_{\underline{b}} H_m(N_f E_{\underline{b}}) \cong \bigoplus_{\underline{b}} H_m(E_{b_0}),$$

where the isomorphism $\cong$ is that induced by μ (cf. 3.1.1). The second sheet of this spectral sequence is obtained by computing the horizontal homology (n -direction). In degree m, n this is equal to $H_n H_m(C_f E) = H_n(B, H_m(F))$. Finally, by the generalized Eilenberg-Zilber theorem (cf. [GJ, IV,2.5]) the homology of the total complex $H_{m+n}(\text{Tot}(C_f E))$ is isomorphic to the homology of the diagonal $H_{m+n}(d(\mathbb{Z}N_f E))$, which equals $H_{m+n}(E)$ since $d(N_f E)$ and NE are homotopic (cf. 4.1.3). This completes the proof. $\square$

Suppose now that $\xi = (p : E \rightarrow B)$ is a fibration whose fibers are homologically trivial, namely $H_m(E_b) = 0$ if $m > 0$ and $H_0(E_b) = \mathbb{Z}$ for all object b of B . If $m > 0$ then the functors $H_m(F)$ are constant and equal to 0, so the second sheet of the spectral sequence X of 5.2.1 is

$$X_{m,n}^2 = \begin{cases} 0 & \text{if } m > 0 \\ H_n(B) & \text{if } m = 0 \end{cases}$$

It follows that $X^\infty = X^2$ and thus the homology of E is that of B . It is not hard to see that $p_* : H_n(E) \rightarrow H_n(B)$ is actually the isomorphism.

5.2.2 Corollary. If a fibration $\xi = (p : E \rightarrow B)$ is such that the fibers E_b are homologically trivial, then $p_* : H_n(E) \xrightarrow{\sim} H_n(B)$ is an isomorphism for all $n \geq 0$.

5.2.3 Corollary (Homology version of Quillen's Theorem A). Let $u : A \rightarrow B$ be a map between small categories whose homotopy fibers A/b are homologically trivial. Then $u_* : H_n(A) \xrightarrow{\sim} H_n(B)$ is an isomorphism for all $n \geq 0$.

Proof. If $\pi : E(u) \rightarrow B$ is the fibration associated to u , then its fibers are isomorphic to the homotopy fibers of u and thus π induces isomorphisms in homology by 5.2.2. Since $u = \pi \circ i$ and $i : A \rightarrow E(u)$ is a weak equivalence (cf. 2.2.1) the result follows. $\square$

5.3 Quillen fibrations and Theorem B

We have seen in many examples that the several fibers of a Grothendieck fibration need not have the same homotopy type. For instance, the Grothendieck construction over a functor $F : B \rightarrow \text{Cat}$ has as fibers the values of F . This remark shows that in general the map $B_f E \rightarrow B_f B$ is not a fibration, nor a quasifibration. It is remarkable that this is the only obstruction, as was pointed out by Quillen [Qu]. In this section we define Quillen fibrations, discuss the monodromy action and reformulate Quillen's Theorem B in terms of the fibred nerve.

We say that a fibration $\xi = (p : E \rightarrow B)$ is a *Quillen fibration* if for each arrow $\phi : b \rightarrow b'$ in B the base-change functor $\phi_* : E_b \rightarrow E_{b'}$ is a weak equivalence. Note that this definition does not depend on the cleavage for two base-change functors over ϕ must be homotopic.

In a Quillen fibration the induced functor $B \rightarrow [\text{Top}]$, $b \mapsto BE_b$ is morphism inverting, therefore it induces a map

$$\pi_1(B) \rightarrow [\text{Top}].$$

Here $\pi_1(B)$ denotes the fundamental groupoid of B , namely the groupoid obtained by formally inverting all the arrows of B , and $[\text{Top}]$ denotes the category of topological spaces and homotopy classes of continuous maps. We call $\pi_1(B) \rightarrow [\text{Top}]$ the *monodromy action* of the fibration. The monodromy action is a first tool to classify Quillen fibrations. In very special situations, it suffices to recover the whole fibration, as we can see in the following example.

Example. (cf. [Qu]) If $p : E \rightarrow B$ is a Quillen fibration with discrete fibers (the only arrows in the fibers are identities), then the base-change functors $E_b \rightarrow E_{b'}$ must be bijections. It follows that a Quillen fibration with discrete fibers is essentially the same as a functor $B \rightarrow \text{Set} \subset \text{Cat}$ which is morphism inverting, or what is the same, a functor $\pi_1(B) \rightarrow \text{Set}$.

A Quillen fibration $p : E \rightarrow B$ with discrete fibers should be thought of as a covering of categories. Indeed, they yield coverings after applying the classifying space functor.

One is interest in understand how fibrations behave with respect to the classifying space functor. Next example shows that $B_f E \rightarrow B_f B$ need not to be a fibration, so we shall look for a notion weaker than that.

Example. Let E be the full subcategory of $I \times I$ with objects $(1, 0)$, $(0, 1)$ and $(1, 1)$. Then the second projection $E \rightarrow I$ is a fibration. Since the fibers are contractible, it is a Quillen fibration indeed. Despite this, the induced map of topological spaces is not a fibration, as one can easily check.

Recall that a *quasifibration* of topological spaces $f : X \rightarrow Y$ is a map such that the inclusions of the actual fibers into the homotopy fibers are a weak homotopy equivalences. They extend the notion of fibration, and their most important feature is that they yield long exact sequences relating the homotopy groups of the fibre, the total space and the base space.

5.3.1 Theorem. If $p : E \rightarrow B$ is a Quillen fibration, then the induced map $p_* : B_f E \rightarrow B_f B$ is a quasifibration of topological spaces.

Proof. It essentially that of [Qu, lemma p.14]. One proves that the restriction of p_* to the n -th skeleton of $B_f B$ is a quasifibration by induction on n , from where the result follows. To prove the inductive step one writes $sk_n(B_f B)$ as the union $U \cup V$, where U is the open set obtained by removing the barycenters of the n -cells and V is the union of the interiors of the n -cells, and prove that p_* is a quasifibration when restricted to U , V and $U \cap V$.

To prove that $p_*|_U$ is a quasifibration one may proceed in the same fashion as in [Qu]. With V and $U \cap V$ one must be careful: $p|_V$ and $p|_{U \cap V}$ are not product fibrations. Anyway, they can be deformed into product fibrations using 3.1.1. $\square$

Last theorem shows an interesting feature of the fibred nerve: it carries Quillen fibrations into quasifibrations. The question of whether or not $BE \rightarrow BB$ is a quasifibration is rather unclear, and this can be understood as a disadvantage of the classic nerve when dealing with fibrations.

5.3.2 Corollary (Quillen's Theorem B). If $u : A \rightarrow B$ is a map between small categories such that $A/b \rightarrow A/b'$ is a weak equivalence for all $b \rightarrow b' \in \text{fl}(B)$, then there is a long exact sequence

$$\dots \xrightarrow{\partial} \pi_k(A/b, \bar{a}) \rightarrow \pi_k(A, a) \xrightarrow{u_*} \pi_k(B, b) \xrightarrow{\partial} \pi_{k-1}(A/b, \bar{a}) \rightarrow \dots$$

where $a \in \text{ob}(A)$, $b = u(a)$ and $\bar{a} = (a, \text{id}_b)$.

Proof. Let $i : A \rightarrow E(u)$ be the canonical map, r its right adjoint and $w : A/b \rightarrow A$ be the map $(a, u(a) \rightarrow b) \mapsto a$. In the diagram

$$\begin{array}{ccccc} A/b & \xrightarrow{w} & A & \xrightarrow{u} & B \\ \parallel & & \uparrow r & & \parallel \\ E(u)_b & \xrightarrow{\hookrightarrow} & E(u) & \xrightarrow{\pi} & B \end{array}$$

the left square commutes, and since $\pi i = u$ and r is a homotopy inverse to i the right square commutes up to homotopy. We conclude that the homotopy groups of A/b , A and B can be identified naturally with that of the base, the fiber and the total category of the associated fibration $E(u) \rightarrow B$. It is a Quillen fibration, and thus the result follows from 5.3.1. $\square$

5.4 Group actions and TCP

Regarding small categories as combinatorial models for homotopy types, it is natural to investigate their equivariant homotopy properties. In this section we derive a splitting fibration from a small category endowed with a group action, and relate its cleaved nerve with a twisted cartesian product in the sense of [Ma]. We also study the spectral sequence 5.2.1 in this particular case.

A simplicial group G operates on a simplicial set K from the left if there is a simplicial map $G \times K \rightarrow K$, $(g, k) \mapsto g \cdot k$ satisfying $1_n \cdot k = k$ for all $k \in K_n$ and $g_1 \cdot (g_2 \cdot k) = (g_1 g_2) \cdot k$ for all $k \in F_n$ and $g_1, g_2 \in G_n$. Here 1_n denotes the unit of G_n . Given A, B simplicial sets and G a simplicial group which operates on A from the left, a *twisted cartesian product* (TCP) with fibre A , base B and group G is a simplicial set $A \times_\tau B$ with simplices $(A \times_\tau B)_n = A_n \times B_n$ and faces and degeneracies given by

$$d_i(a, b) = (d_i a, d_i b), \quad i > 0; \quad d_0(a, b) = (\tau(b) \cdot d_0 a, d_0 b), \quad \tau(b) \in G; \quad s_i(a, b) = (s_i a, s_i b), \quad i \geq 0.$$

Here $\tau : B_n \rightarrow G_{n-1}$ is a function which must satisfy some standard identities in order to make $A \times_\tau B$ a simplicial set. This τ is called the *twisting function*.

Let G be a group, and let A be a small category on which G acts. This action can be seen as a group morphism $G \rightarrow \text{Aut}(A)$, $g \mapsto u_g$, or equivalently, as a functor $G \rightarrow \text{Cat}$ that maps the unique

object of G to $A \in \text{ob}(\text{Cat})$. The Grothendieck construction over this functor is a splitting fibration $p : G \rtimes A \rightarrow G$ over G .

The constant simplicial group G (on which every face and degeneracy operator is the identity) operates on NA from the left via the formula

$$g \cdot (a_0 \rightarrow \dots \rightarrow a_n) = (u_g(a_0) \rightarrow \dots \rightarrow u_g(a_n)).$$

5.4.1 Proposition. The diagonal of the cleaved nerve $dN_c(G \rtimes A)$ can be regarded as a TCP between the nerves of A and G , namely $NA \times_\tau NG$.

Proof. Let $\tau : NG_n \rightarrow G_{n-1} = G$ be the projection $\tau(* \xrightarrow{g_1} * \xrightarrow{g_2} \dots \xrightarrow{g_n} *) = g_1$, and let $NA \times_\tau NG$ be the TCP with twisting function τ . We define a simplicial map $\varphi : dN_c(G \rtimes A) \rightarrow NA \times_\tau NG$ by giving to each simplex $s \in N_c(G \rtimes A)_{n,n}$ the pair (a, b) , where a is the mast of s and b is its base. One checks easily that φ is actually a simplicial map, and it is an isomorphism because $G \rtimes A \rightarrow G$ is splitting and 3.2.1. $\square$

Given G acting over A , the fibration $G \rtimes A \rightarrow G$ has a unique fiber, which is isomorphic to A . Thus the modules $H_m(F)$ of the spectral sequence 5.2.1 are just the homology groups of A endowed with the action of G . Writing $A//G = G \rtimes A$ for the homotopy theoretic quotient (cf. 5.1.2) we obtain the following version of the Eilenberg-Moore spectral sequence (cf. [An, p.775]) as an application of 5.2.1.

5.4.2 Proposition. There is a spectral sequence $\{X_{m,n}^r\}$ which converges to the homology of the homotopy theoretic quotient $A//G$ and whose second sheet consists of the group homology of G with coefficients in the homology of A .

$$X_{m,n}^2 = H_n(G, H_m(A)) \Rightarrow H_{m+n}(A//G)$$

References

- [An] D. Anderson. *Fibrations and geometric realizations*. Bull. of the Am. Math. Soc. 84 (1978).
- [Bo] F. Borceux. *Handbook of Categorical Algebra 2, Categories and Structures*. Encyclopedia of Math and its App 51, Cambridge (1994).
- [BK] A.K. Bousfield, D.M. Kan. *Homotopy Limits, Completions and Localizations*. Lecture Notes in Math. vol. 304, Springer (1972).
- [CR] A. Cegarra, J. Remedios. *The relationship between the diagonal and the bar constructions on a bisimplicial set*. Topology and its App 153 (2005), 21-51.
- [dH] M. del Hoyo. *On the subdivision of small categories*. Topology and its App 155 (2008), 1189-1200.
- [Ev] M. Evrard. *Fibrations de petites categories*. Bull. Soc. math. France 103 (1975), 241-265.
- [GJ] P. Goerss, J. Jardine. *Simplicial homotopy theory*. Progress in Math 174. Birkhuser Verlag (1999).
- [SGA1] A. Grothendieck. *Revetements etales et groupe fondamental (SGA 1)*. Springer Lecture Notes in Math., Vol. 224.
- [He] M. Heggie. *Homotopy cofibrations in Cat*. Cahiers Topologie Geom Diff 33 (1992), 291-313.
- [Il] L. Illusie. *Complexe cotangente et deformations II*. Lecture Notes in Math 283, Springer (1972).
- [Ma] J. P. May. *Simplicial objects in algebraic topology*. Van Nostrand Mathematical Studies 11 (1967).

- [Mi] G. Minian. *Numerably Contractible Categories*. K-Theory 36 (2005), 209–222.
- [Qu] D. Quillen. *Higher Algebraic K-Theory I*. Lecture Notes in Math. 341, Springer (1973), p. 85–147.
- [Se] G. Segal. *Classifying space and spectral sequences*. Pub. I.H.E.S. 34 (1968), 105–112.
- [Th] R. Thomason. *Homotopy colimits in the category of small categories*. Math. Proc. Camb. Phil. Soc. 85 (1979), 91–109.